

# Spacewatch and Follow-up Astrometry of Near-Earth Objects

International Asteroid Warning Network Steering Group  
Meeting

Cambridge, MA 2014 Jan 13

Robert S. McMillan<sup>1</sup>, T. H. Bressi<sup>1</sup>, J. A.  
Larsen<sup>2</sup>, J. V. Scotti<sup>1</sup>, and R. A. Mastaler<sup>1</sup>

URL: <http://spacewatch.lpl.arizona.edu>

<sup>1</sup>University of Arizona; <sup>2</sup>U.S. Naval Academy

# Summary

- Follow-up of "large" NEOs ( $H \leq 22$ ) as they recede from Earth after discovery and become fainter, as well as VIs, PHAs, & NEOs observed by WISE.
- New, fast-reading CCD on 1.8-meter telescope.
- Observed at elongations as small as  $46^\circ$ .
- ~2800 tracklets of NEOs accepted by MPC from Spacewatch each year.
- Big, long archive from 0.9-m telescope to support precoveries.

# Why Targeted Followup is Needed

- Discovery arcs too short to define orbits:
  - Followup observation intervals need to be thousands of times longer than discoveries.
- Objects can escape redetection by surveys:
  - Surveys too busy covering other sky.
  - Objects tend to get fainter after discovery.
- Sky density of detectable NEOs is too sparse to rely on incidental redetections alone.

# Why More Followup is Needed

- 1/3rd of PHAs observed on only 1 opposition.
- 1/6th of PHAs' arcs  $< 30^{\text{d}}$ .
- ~Half of potential close approaches in the next 30 years will be by objects observed on only one opposition.
- 2/3<sup>rds</sup> of  $H \leq 22$  VI's on JPL risk page *are lost* and  $>$  half of those were discovered within the last 6 years by modern surveys.

# How “lost” can they get?

- (719) Albert discovered visually in 1911.
- “Big” Amor asteroid, diameter  $\sim 2$  km.
- Favorable (perihelic) apparitions 30 yrs apart.
- Missed in 1941 due to inattention.
- Missed in 1971 due to large uncertainty.
- MPC recognized (719) as a rediscovery by Spacewatch in 2000.

# 1979 XB: A “Big” Lost “VI”!


- 4-day observed arc in 1979 December.
- $H \approx 18.5 \leftrightarrow$  Diameter 370-1200 m.
- Synodic period  $\approx 1.4^y$ .
- Possible close encounters in 2056 & 2086.
- Not rediscovered in  $>3$  decades of modern surveying.

# 0.9-m Telescope Modernized by Spacewatch in 2002

- Hyperboloidal primary & refractive field corrector.
- Mosaic of 4 CCDs.
- Bandpass  $\approx 0.5\text{-}0.9 \mu\text{m}$ ;  $\lambda_{\text{eff}} \approx 0.7 \mu\text{m}$ .
- Began 2003 April; 22 nights per lunation.
- Automated in 2005 May.
- Patterns near opp'n, & low elongation in east.
- $1400 \text{ deg}^2$  per lunation;  $V \text{ mag} \approx 20.5\text{-}21.7$ .

# 0.9-m Telescope in 2012

Photo by Roger Carpenter, 2012 Feb






## Spacewatch CCD Mosaic on 0.9-m telescope.

Four EEV Grade-1,  
back-illuminated,  
antireflection-coated  
CCDs of 4608x2048  
pixels each.

37 million pixels.

1 arcsec per pixel.

2.9 deg<sup>2</sup> covered.



# Archive from Mosaic on 0.9-m:

- Revisits @ 4<sup>d</sup> intervals aid MBA linkages.
- ~20 TB in size.
- 11 yrs of uniformly conducted surveying.
- Incidental astrometry & precoveries of NEOs.
- V mag limit ~20-21.
- Coverage ~1400 deg<sup>2</sup> per lunation (3 passes per pointing) mostly along ecliptic and low-elongation in the east.

## Spacewatch 1.8-meter Telescope on Kitt Peak

**New CCD in 2011 Oct:**

FOV =  $20' \times 20'$ .

Scale = 0.6 arcsec/pixel.

Bandpass  $\square$  “V+R+I”.

Fast readout.

Limit V=23.

**54% more obs per year.**

Astrometric resids  $\square$  **0.3  
arcsec**, vs. 0.6 on NEOs  
with the old CCD.

Photo by Roger Carpenter, 2012 Feb.



## 2.3-meter Bok Telescope of Steward Observatory on Kitt Peak

90Prime mosaic camera:

FOV  $\sim 1 \text{ deg}^2$

0.45"/pixel.

V mag limit  $\approx 24$ .

$\sim 24$  nights per year.

$\sim 3\text{-}4$  objects per hour.

Photo 2007 by Marc Murison, USNOFS.



## 4-meter Mayall Telescope of Kitt Peak National Observatory


Prime focus mosaic of  
CCDs:

FOV= 37' x37'.

Time awarded to  
Spacewatch for faint  
( $V \geq 23$ ) priority NEO  
followup.

~150 object-visits per yr.


Photo: NOAO/AURA/NSF



**3.5-m telescope of  
Wisconsin-Indiana-  
NOAO (WIYN) on Kitt  
Peak, Az.**

Used by Spacewatch  
in 2010 in Target-  
of\_Opportunity (ToO)  
mode to recover  
selected faint NEOs  
discovered by the  
Near-Earth Object  
Wide-field Infrared  
Survey (NEOWISE)  
spacecraft mission.


Photo: NOAO/AURA/NSF; copyright  
WIYN Consortium, Inc., all rights  
reserved.



# Spacewatch Contributions

- Between 2011 Oct 16 and 2013 Dec 5  
Spacewatch observed:
  - 53% of all NEOs observed in that time.
  - 59% of all PHAs observed in that time.
- Leading station in followup of provisionally designated PHAs while faint ( $V \geq 21.5$ ); contributing 36% of all such observations.

# Cumulative Spacewatch Observations of PHAs





# Spacewatch Observations of WISE-detected Asteroids

- Recoveries & astrometry improve orbits.
- Photometry supports albedo determination.
- Lightcurves reveal rotation period, amplitude, & and rotational phase.
- *BVRIZ* taxonomic photometry to compare with albedos & orbital classes.

# Needs of Followup Campaign (besides money).

- Get longer arcs during discovery apparitions:
  - Keep provisional designations on the NEO CP.
  - Encourage more time on larger telescopes.
- Software tools to grade followup stations.
- MPES-format-compatible lists of targets on NHATS, NEOWISE, & Radar websites.
- More reliable NEOCP scores & uncertainty maps.
- More focused selections of targets in NEA Observation Planner.
- More followup stations blogging targets in advance.

# Acknowledgements

- NASA's NEO Observation Program.
- The IAU's Minor Planet Center.
- JPL's NEOWISE Team led by A. K. Mainzer.
- JPL's NEO Office.
- Kitt Peak Nat'l & Steward Observatories.
- The Brinson Foundation of Chicago, IL.
- The U. S. Naval Academy.
- The estates of R. L. Waland and R. S. Vail.
- Other private donors.